

XXI. NWA 1068 and NWA 1110 (ver. 2003)

basalt, ~700 grams
find



Figure XXI-1: Photograph of NWA 1068 (scale is 1 cm). Photo kindly provided by Bruno Fectay and Carine Bidaut.

Introduction

NWA 1068 was found in the Moroccan Sahara in April 2001 by a local team called “La Mémoire de la Terre” and given the field name “Louise Michel”. It has no fusion crust and is greenish-brown in color (figure XXI-1). A main mass (522g) and 22 small fragments were recovered (Barrat *et al.* 2002b). NWA 1110 (several fragments, 118 grams) was purchased from meteorite dealers in Morocco in September 2001 and appears to be paired with NWA 1068 (Irving and Kuehner). The exact strewn field of this meteorite is not known, but appears to be in Morocco (Barrat *et al.* 2002; Russell *et al.* 2002).

In texture, mineralogy and bulk chemistry, NWA 1068 appears similar to other basaltic shergottites,

EETA79001, Dar al Gani 476, Sayh al Uhaymir 005 and Dhofar 019, but the REE pattern of NWA 1068 is not depleted in light REE and is instead very similar to that of Shergotty.

Petrography

NWA 1068 has a porphyritic texture consisting of olivine phenocrysts (up to 2 mm) in a fine-grained groundmass (grain size ~100 microns). The olivine phenocrysts occur in clusters, contain chromite and melt inclusions and appear corroded/resorbed (Barrat *et al.* 2002).

Wadhwa and Crozaz (2002) analyzed a melt inclusion with an olivine, and found that it had a REE pattern parallel to that of the bulk rock, indicating that the

Mineralogical Mode

	Barrat <i>et al.</i> (2002a, b)	Mikouchi (2002)
Olivine	21 vol. %	27.2 %
Pyroxene	52	51.6
Maskelynite	22	15.5
Phosphate	2	1.8
Opaques	2	2.5
Mesostasis	1	

olivine phenocrysts formed from a magma similar to that of the NWA 1068 parent melt.

NWA 1068 contains veins of terrestrial calcium carbonate, small “melt” pockets and numerous shock veins (Barrat *et al.* 2002b).

Mineral Chemistry

Olivine: The cores of the olivine megacrysts are homogeneous (Fo₇₀) with distinct outer rims up to 100 microns wide, zoned to Fo₅₀₋₆₀. There is no outer zoning where olivine grains are touching.

Pyroxene: Both pigeonite and augite are present as subhedral to euhedral crystals in about equal proportions (figure XXI-2). Pyroxenes are normally zoned (En₅₇Wo₅ to En₄₀Wo₁₃ and En₅₅Wo₂₁ to En₃₅Wo₂₈).




Figure XXI-2: Pyroxene and olivine composition diagram for NWA 1068 (data replotted from Barret *et al.* 2002).

Maskelynite: Shocked plagioclase crystals are normally zoned An₅₃ to An₄₉.

Opaque minerals: Chromite is found associated with olivine. Ilmenite and ulvöspinel are found in the mesostasis.

Phosphates: Barrat *et al.* (2002) reported analyses of both merrillite and apatite in NWA 1068.

Whole-rock Composition

NWA 1068 has a relatively flat REE pattern, unlike that of the other olivine-bearing shergottites, which have (rather extreme) light REE depletions (figure XXI-3). Key element ratios such as Fe/Mn (= 45), Al/Ti




Figure XXI-3: Rare earth element diagram for NWA 1068 compared with Shergotty (data from Barret *et al.* 2002 and Lodders 1998).

(=6.6) and Ga/Al (= 4.4 x 10⁻⁴) are typical of Martian meteorites.

Sr, Ba and Pb are relatively high compared with unweathered shergottites (figure XXI-4), but Th/U is normal and NWA 1068 does not have a Ce anomaly, so it has been concluded that this “hot desert” find is not badly weathered (Barrat *et al.* 2002b).

Radiogenic Isotopes

Shih *et al.* (2003) have dated NWA 1068 by Rb-Sr and Sm-Nd (figures XXI-6 and XXI-7). The apparent crystallization age (185 Ma) is typical of basaltic shergottites.

Cosmogenic Isotopes

Other Isotopes

Shock Effects

Thin shock veins and small pockets of shock melt glass are found in the groundmass. Barrat *et al.* (2002) report analyses of this glass.

Terrestrial Weathering

The occurrence of veins of calcium carbonate demonstrates that NWA 1068 has been weathered after its fall. It has elevated Sr, Ba and Pb concentrations

(Barrat *et al.* 2002). Surfaces are coated with reddish desert varnish.

Processing Info

The sizes of the small pieces collected along with NWA 1068 referred to above are (19.2 g, 10.4 g, 6.0 g, 3.2 g, 2.0 g, 2.4 g, 1.9 g, 1 g, 1 g, 0.9 g, 0.8 g, 0.6 g, 0.5 g, 0.4 g, 0.3 g etc). NWA 1110 was also in pieces, and there are certain to be others, still to be found !




Figure XXI-4: Trace element data for basaltic shergottites normalized by composition of Shergotty (average by Lodders 1998).



Figure XXI-5: Photograph of pieces of NWA 1110 (probably paired with NWA 1068). Photograph and information from JPL web site (Ron Baalke).

Table XXI-1: Composition of NWA1068.

reference	Barrat 2002	
weight		
SiO2	0.77	(a)
TiO2	5.75	(a)
Al2O3	20.48	(a)
FeO	0.46	(a)
MnO	7.91	(a)
CaO	16.5	(a)
MgO	1.14	(a)
Na2O	0.16	(a)
K2O		
P2O5		
sum		
Li ppm	4.34	(b)
Be	0.35	(b)
Sc	37	(b)
V	280	(b)
Cr	4317	(b)
Co	56.2	(b)
Ni	232	(b)
Cu	14	(b)
Zn	49	(b)
Ga	13.4	(b)
Ge		
Rb	5.75	(b)
Sr	67	(b)
Y	17.19	(b)
Zr	62.14	(b)
Nb	4.37	(b)
Mo		
Cs ppm	0.45	(b)
Ba	127	(b)
La	2.25	(b)
Ce	5.38	(b)
Pr	0.783	(b)
Nd	3.82	(b)
Sm	1.49	(b)
Eu	0.552	(b)
Gd	2.14	(b)
Tb	0.414	(b)
Dy	2.8	(b)
Ho	0.59	(b)
Er	1.63	(b)
Tm		
Yb	1.37	(b)
Lu	0.198	(b)
Hf	1.58	(b)
Ta	0.2	(b)
W ppb	520	(b)
Th ppm	0.409	(b)
U ppm	0.1	(b)

technique: (a) ICP/AES, b) ICP-MS




Figure XXI-6. Rb-Sr isochron from Shih et al. 2003.




Figure XXI-7. Sm-Nd isochron for NWA1068 by Shih et al. 2003.